
ALDEHYDE, KETONE, CARBOXYLIC ACID :-

(ORGANIC CHAPTER-3)


The functional group >C=O is called carbonyl group. Organic compounds containing carbonyl groups are aldehydes, ketones, carboxylic acids and their derivatives. The general formulae of these compounds are given below:-



Aldehyde

IUPAC :- Alkanal


(where R may be H or any alkyl, aryl or aralkyl group)



Ketone

IUPAC :- Alkanone

(where R and R' may be same or different alkyl, aryl or aralkyl group)




Carboxylic acid

IUPAC :- Alkanoic acid


(where R may be H or any alkyl, aryl or aralkyl group)

Derivatives of Carboxylic acids :-




Acyl halide

(X = halogen)




Acid anhydride

(where R & R' may be same or different groups)



Ester














(where R & R' may be same or different groups)



Acid amide

(where R may be H or any group)

IUPAC names of some aldehyde & ketone that were asked in the board examination :-

	STRUCTURE	IUPAC name	STRUCTURE	IUPAC name
1.		Propanal	9. 	Pentane-2,4 dione
2.		4-hydroxypentan-2-one	10. 	But-3-en-2-one
3.		2-methylpropan-1-al	11. 	2-hydroxybenzal dehyde.
4.		Pent-2-en-1-al	12. 	3-oxopentanal
5.		4-methylpent-3-en-2-one	13. 	Hex-2-en-4-ynoic acid
6.		4-chloropentan-2-one		
7.		2-chlorophenylethanone		
8.		4,4'-dihydroxy benzophenone.	14. 	Benzene-1,4-di carboxylic acid.

PREPARATION OF ALDEHYDE AND KETONE:-

① Oxidation of alcohol

1° Alcohol + PCC ⟶ Aldehyde $\xrightarrow{PCC}$ No product

1° Alcohol + H_2SO_4 $\xrightarrow{K_2Cr_2O_7}$ Aldehyde $\xrightarrow[K_2Cr_2O_7]{H_2SO_4}$ Carboxylic Acid

Eg:- $CH_3CH_2OH \xrightarrow{PCC} CH_3CHO$

Ethanol → Ethanal


$CH_3CH_2OH \xrightarrow[H_2SO_4]{K_2Cr_2O_7} CH_3CHO \xrightarrow[K_2Cr_2O_7]{H_2SO_4} CH_3COOH$

Ethanol → Ethanal → Ethanoic acid

2° Alcohol $\xrightarrow{PCC}$ Ketone

2° Alcohol $\xrightarrow[H_2SO_4]{K_2Cr_2O_7}$ Ketone

Eg:-



Cyclohexanone

$CH_3-\underset{\underset{OH}{|}}{CH}-CH_3 \xrightarrow{PCC} CH_3-\overset{\overset{O}{||}}{C}-CH_3$

N.B:- 3° Alcohol $\xrightarrow{PCC}$ No product

3° Alcohol never undergo oxidation because it has no α-H atom

② Dehydrogenation of alcohols:-

1° Alcohol $\xrightarrow{Cu/573K}$ Aldehyde + H_2

Eg:- $CH_3-CH_2-OH \xrightarrow{Cu/573K} CH_3CHO + H_2$

2° Alcohol $\xrightarrow{Cu/573K}$ Ketone + H_2

Eg:- $CH_3-\underset{\underset{OH}{|}}{CH}-CH_3 \xrightarrow{Cu/573K} CH_3-\overset{\overset{O}{||}}{C}-CH_3 + H_2$

3° Alcohol $\xrightarrow{Cu/573K}$ Alkene + H_2O

Eg:- $CH_3-\underset{\underset{CH_3}{|}}{\overset{\overset{CH_3}{|}}{C}}-OH \xrightarrow{Cu/573K} CH_3-\underset{\underset{CH_3}{|}}{C}=CH_2 + H_2O$


③ FROM HYDROCARBONS:-

(a) Oxonolysis of alkenes:-


Alkene	$\xrightarrow[Zn/H_2O]{O_3}$	2 Aldehyde / 2 Ketone / 1 Aldehyde or 1 Ketone
$CH=CH$	$\xrightarrow[Zn/H_2O]{O_3}$	2 Aldehyde
$C=CH$	$\xrightarrow[Zn/H_2O]{O_3}$	1 Aldehyde + 1 Ketone
$C=C$	$\xrightarrow[Zn/H_2O]{O_3}$	2 Ketone

Eg:- $CH_3-CH=CH_2 + O_3 \xrightarrow{Zn/H_2O} CH_3CHO + H-\overset{O}{\overset{\|}{C}}-H$

Propene → Ethanal + Methanal



Benzaldehyde



Acetone


TRICK:- Remove the double bond and provide it to each C-atom.

PREPARATION OF ALDEHYDE:-


① FROM ACID CHLORIDE (ROSENMOND REDUCTION REACTION)

When H_2 gas is passed over acid chloride in presence of Pd supported with $BaSO_4$ then partially poisoned by sulphur or quinoline forms aldehyde.

$$\text{Acid chloride} \xrightarrow[Pd-BaSO_4]{H_2} RCHO + HCl$$



Eg:- $CH_3-\overset{O}{\overset{\|}{C}}-Cl \xrightarrow[Pd-BaSO_4]{H_2} CH_3-\overset{O}{\overset{\|}{C}}-H + HCl$



Benzoyl chloride → Benzaldehyde


② FROM NITRILES (STEPHEN REACTION)

$$R-C\equiv N \xrightarrow[H_2]{SnCl_2/HCl} \underset{\text{Imine}}{R-CH=NH} \xrightarrow{H_3O^{\oplus}} RCHO$$

Eg:-

$CH_3CN \xrightarrow[H_2]{SnCl_2/HCl} CH_3CH=NH \xrightarrow{H_3O^+} CH_3CHO$

Acetonitrile → Acetaldehyde



Benzonitrile → Benzaldehyde

TRICK:- When imine is prepared, one H is attached to carbon and another H is attached to Nitrogen.

(3) Using DIBAL-H (di-isobutyl aluminium hydride)

$$R-C\equiv N \xrightarrow[H_2O]{DIBAL-H} R-CHO$$

Eg:- $CH_3-CH=CH-CN \xrightarrow[H_2O]{DIBAL-H} CH_3-CH=CH-CHO$

(4) From ester:-

$$R-\overset{O}{\overset{\|}{C}}-O-R' \xrightarrow[H_2O]{DIBAL-H} R'OH + R-\overset{O}{\overset{\|}{C}}-H$$

Eg:- $CH_3-CH_2-\overset{O}{\overset{\|}{C}}-O-C_2H_5 \xrightarrow[H_2O]{DIBAL-H} C_2H_5OH + CH_3-CH_2-\overset{O}{\overset{\|}{C}}-H$


TRICK:- the bond adjacent to C=O breaks and H is attached to each 1 of the group.

(5) FROM HYDROCARBONS:-

(i) By oxidation of methyl benzene:-


(a) Use of chromyl chloride (ETARD REACTION)

When Toulene is treated with CrO_2Cl_2 in presence of non-polar solvent like CS_2 then it forms a chromium complex, which on further hydrolysis forms aldehyde.




Toulene → Chromium complex → Benzaldehyde

(b) USE OF CHROMIC OXIDE:-




Toulene — Benzaldehyde

(ii) Side chain chlorination followed by hydrolysis:-



Toulene — Benzal chloride — Benzaldehyde


(iii) By Gatterman-Koch reaction:-




Benzene — Benzaldehyde

PREPARATION OF KETONES:-


① FROM ACID CHLORIDES:-




Eg:- 


(Propionyl chloride)

② FROM NITRILES:-




Eg:- 

③ FRIEDEL-CRAFTS ACYLATION:-



Benzene — Acetophenone

Eg:- 

Benzene — Propiophenone

PHYSICAL PROPERTIES:-

① BOILING POINT:-

Carboxylic acid > Alcohol > Aldehyde > Ketone > Ester > Hydrocarbon

The boiling points of aldehydes and ketones are higher than hydrocarbons and ethers of comparable molecular masses. It is due to weak molecular association in aldehydes and ketones arising out of the dipole-dipole interactions.


② SOLUBILITY:-

* Lower aldehyde and ketone are soluble in water because of intermolecular hydrogen bonding.
* Higher aldehyde and ketone are insoluble in water.

CHEMICAL PROPERTIES:-

1. Aldehyde and ketone undergo nucleophilic addition reaction.

(i) Mechanism of nucleophilic addition reaction:-



The sp^2 hybridised carbon atom changes to sp^3 hybridised carbon atom.

(ii) Reactivity:-

Aldehyde are more reactive than ketone because:-

① In case of ketone, due to presence of more +I group, the electrophilicity of carbonyl carbon atom decreases and hence, ketone is less reactive towards nucleophilic addition reaction.


Eg:- $CH_3-\overset{O}{\overset{\|}{C}}-H > CH_3-CH_2-\overset{O}{\overset{\|}{C}}-H > CH_3-\underset{CH_3}{\overset{CH_3}{\underset{|}{\overset{|}{C}}}}-\overset{O}{\overset{\|}{C}}-H$

② Steric factor:- In case of ketone, due to presence of 2 bulkier alkyl group, the approach of nucleophile is hindered due to steric repulsion.

(iii) Some important examples of nucleophilic addition and nucleophilic addition-elimination reactions:-


(a) Addition of HCN:-

Since, HCN is a weak acid, it generates nucleophile in basic medium




Cyanohydrin complex

Eg:-




Cyanohydrin complex




Cyanohydrin complex

(b) Addition of sodium hydrogen sulphite:-



(c) Addition of alcohol:-


Aldehyde react with monohydric alcohol to form hemiacetal and acetal.



Hemi-acetal

Acetal

Eg:-




Hemiacetal


Acetal

* Reaction with ethylene glycol :-

Ketone react with dihydric alcohol to form ketals.




Eg :-



Acetone — Ethylene glycol — Ethylene glycol ketal.


(d) Addition of ammonia and its derivatives :-

① Reaction with ammonia :- (NH_3)




Imine

Eg :-




Imine

② Reaction with amine :- ($R-NH_2$)




Substituted Imine

Eg :-




③ Reaction with hydroxyl amine (NH_2OH) :-




Oxime complex

Eg :-




Ethanal — Ethanal oxime

④ Reaction with hydrazine (NH_2-NH_2):-



Hydrazone


Eg:-



Acetaldehyde


Acetaldehyde hydrazone

⑤ Reaction with phenylhydrazine ($C_6H_5-NH-NH_2$):-



Phenyl hydrazone

Eg:-




Acetaldehyde

Acetaldehyde phenylhydrazone

⑥ Reaction with 2,4 DNP (2,4-dinitro phenyl hydrazine):-


* It is used to test the presence of carbonyl group. It gives a orange/red precipitate.



2,4 DNP

2,4-dinitro phenyl hydrazone (Orange or red ppt)

Eg:-




Acetaldehyde

Acetaldehyde 2,4-dinitrophenyl hydrazone.


⑦ Reaction with semicarbazide ($NH_2-NH-\overset{O}{\overset{\|}{C}}-NH_2$):-

* Out of the two NH_2 groups present in semicarbazide, the one which is away from $-\overset{O}{\overset{\|}{C}}-$ is involved in the reaction.



Semicarbazone

Eq:-




Acetaldehyde

Acetaldehyde semicarbazone


2. REDUCTION REACTIONS:-

(i) Reduction to hydrocarbons:-


(a) Clemmensen's reduction:-



Eq:-




(b) Wolff-Kishner reduction:-



Hydrazone


Eq:-




Acetone

3. OXIDATION REACTIONS:-


(a) Oxidation with aldehyde:-



Eq:-



(a) Oxidation with ketone:-



Ketones in presence of conc. HNO_3 (drastic condition) oxidises.

Ketone $\xrightarrow[\Delta]{\text{conc. } HNO_3}$ Two carboxylic acid

In case of unsymmetrical ketones such as Butan-2-one or Pentan-2-one etc, the point of cleavage is such that the keto group stays with smaller alkyl group preferentially. (Popoff's rule)

Eg:-

$CH_3-\overset{\overset{O}{\|}}{C}-CH_2-CH_3 \xrightarrow[\text{conc. } HNO_3]{[O]} CH_3-\overset{\overset{O}{\|}}{C}-OH + CH_3COOH$

Butan-2-one — Ethanoic acid

$CH_3-\overset{\overset{O}{\|}}{C}-CH_2CH_2CH_3 \xrightarrow[\text{conc. } HNO_3]{[O]} CH_3-\overset{\overset{O}{\|}}{C}-OH + CH_3CH_2COOH$

Pentan-2-one — Acetic acid — Propanoic acid

However, in case of symmetrical ketones, only one type of cleavage occurs.

Eg:-

$CH_3CH_2-\overset{\overset{O}{\|}}{C}-CH_2CH_3 \xrightarrow[\text{conc. } HNO_3]{[O]} CH_3CH_2-\overset{\overset{O}{\|}}{C}-OH + CH_3COOH$

Pentan-3-one — Propanoic acid — Ethanoic acid

The mild oxidising agents given below are used to distinguish aldehydes from ketones:-

(i) Oxidation by Tollens' reagent:-

* Tollens' reagent is ammonical silver nitrate solution. $[Ag(NH_3)_2]^+ + OH^-$
* It is the confirmatory test of aldehyde.

$$RCHO + [Ag(NH_3)_2]^+ + OH^- \longrightarrow RCOO^- + Ag + NH_3$$

(Silver mirror)

Eg:- $CH_3CHO + [Ag(NH_3)_2]^+ + OH^- \longrightarrow CH_3COO^- + Ag + NH_3$

N.B:- Ketone $+ [Ag(NH_3)_2]^+ + OH^- \longrightarrow$ No silver mirror (No reaction)

(ii) Oxidation by Fehling's solution:-

Fehling's solution $\longrightarrow$ Fehling A + Fehling B

Fehling A ↓ (aq $CuSO_4$)

Fehling B ↓ (Alkaline sodium potassium tartrate i.e Rochelle salt)

$$RCHO + Cu^{+2} + OH^- \longrightarrow RCOO^- + Cu_2O + H_2O$$

(Red brown ppt)

Eg:-

$$CH_3CHO + Cu^{+2} + OH^- \longrightarrow CH_3COO^- + Cu_2O + H_2O$$

N.B:- Aromatic aldehyde, (benzaldehyde) however doesn't reduce Fehling's solution.

(iii) Iodoform test :-

* It is a chemical test to distinguish the presence of $CH_3-\overset{O}{\overset{\|}{C}}$ group or methyl ketone group in aldehydes and ketones.

$CH_3-CH_2-\overset{O}{\overset{\|}{C}}-H \longrightarrow$ doesn't give iodoform test

$CH_3-\overset{O}{\overset{\|}{C}}-H \longrightarrow$ gives iodoform test.

$$R-\overset{O}{\overset{\|}{C}}-CH_3 + NaOH + I_2 \longrightarrow CHI_3 + R-\overset{O}{\overset{\|}{C}}-ONa$$

(Iodoform)

Eg:- $CH_3-\overset{O}{\overset{\|}{C}}-CH_3 + NaOH + I_2 \longrightarrow CHI_3 + CH_3-\overset{O}{\overset{\|}{C}}-ONa$

4. Reactions due to α-hydrogen:- (ALDOL CONDENSATION)

Aldol is defined as the combination of both alcohol and aldehyde group in two consecutive carbon atoms.

* The aldehyde or ketone must have one α-H atom

(i) Self aldol condensation (If both are same type of reactant):-

$CH_3-\overset{H}{\underset{|}{C}}=O$ + $H-\overset{\alpha}{C}H_2CHO$ $\underset{}{\overset{dil\ NaOH}{\rightleftharpoons}}$ $CH_3-\overset{H}{\underset{OH}{\underset{|}{\overset{|}{C}}}}-CH_2CHO$ $\xrightarrow[-H_2O]{\Delta}$ $CH_3-CH=CH-\overset{O}{\overset{\|}{C}}-H$

Ethanal — 3-Hydroxybutanal — But-2-enal

$CH_3-\overset{CH_3}{\underset{|}{C}}=O$ + $H-CH_2-\overset{O}{\overset{\|}{C}}-CH_3$ $\underset{}{\overset{dil\ NaOH}{\rightleftharpoons}}$ $CH_3-\overset{CH_3}{\underset{OH}{\underset{|}{\overset{|}{C}}}}-CH_2-\overset{O}{\overset{\|}{C}}-CH_3$ $\xrightarrow[-H_2O]{\Delta}$ $CH_3-\underset{CH_3}{\underset{|}{C}}=CH-\overset{O}{\overset{\|}{C}}-CH_3$

Acetone — Diacetone alcohol (aldol) — 4-Methylpent-3-en-2-one


(ii) Cross aldol condensation :-

When 2 different aldehyde both having α-H atom reacts with dil NaOH we get 4 Aldol i.e (A→A, B→B, A→B, B→A)


CH_3CHO + CH_3CH_2CHO ⟶

A B


(A-A condensation)




(B-B condensation)



(A-B condensation)



(B-A condensation)





* In this, we get 2 self aldol products and 2 cross aldol products.

5. OTHER REACTIONS:-

(i) CANNIZZARO REACTION:-

Aldehyde which donot have α-H atom, undergo self oxidation and reduction in presence of conc. NaOH which is reduced to alcohol and oxidised to acid.






Benzyl alcohol Sodium benzoate.





TRICK:- In Cannizzaro reaction, 1 CHO group becomes CH_2OH and another CHO group becomes COONa.

(ii) Electrophilic substitution reaction:-




CARBOXYLIC ACIDS:-

IUPAC name of some carboxylic acid that was asked in the board exam:-

STRUCTURE	IUPAC name
1. $CH_3-CH(OH)-CH_2-COOH$	3-hydroxybutanoic acid
2. 	2-hydroxybenzoic acid
3. $COOH-(CH_2)_4-COOH$	Hexane-1,6-dioic acid
4. $CH_3-C\equiv C-CH=CH-C(=O)-OH$	Hex-2-en-4-yn-oic acid
5. 	3-bromo-5-chlorobenzoic acid
6. 	3,5-dimethylphenyl ethanoate.
7. 	Ethyl-4-chlorobenzoate

STRUCTURE OF CARBOXYL GROUP:-

The carboxylic carbon is less electrophilic than carbonyl carbon atom because of the possible resonance structure:-



PREPARATION OF CARBOXYLIC ACID :-

1. FROM PRIMARY ALCOHOL AND ALDEHYDE :-

$$R-CH_2-OH \xrightarrow[H^{\oplus}]{K_2Cr_2O_7} RCOOH$$

(1° Alcohol)

$$RCHO \xrightarrow[H^{\oplus}]{K_2Cr_2O_7} RCOOH$$


Eg :- $CH_3-CH_2-OH \xrightarrow[H^{\oplus}]{K_2Cr_2O_7} CH_3COOH$


Ethanol — Acetic acid

$$CH_3CHO \xrightarrow[H^{\oplus}]{K_2Cr_2O_7} CH_3COOH$$

Acetaldehyde — Acetic acid

2. FROM ALKYL BENZENES :-





TRICK :- It will convert total side chain to COOH.

3. FROM NITRILES :-


$$R-C\equiv N \xrightarrow[\Delta]{H_3O^{\oplus}} RCOOH + NH_3$$

Eg :- $CH_3-C\equiv N \xrightarrow[\Delta]{H_3O^{\oplus}} CH_3COOH + NH_3$

4. FROM AMIDES :-

$$R-\overset{\overset{O}{\|}}{C}-NH_2 \xrightarrow[\Delta]{H_3O^{\oplus}} RCOOH + NH_3$$

Eg :-



Benzamide — Benzoic acid

5. FROM GRIGNARD REAGENT :-

* It is the method to increase one C.

$$R-MgBr \xrightarrow[(ii)\ H_3O^{\oplus}]{(i)\ CO_2} RCOOH$$

Eg:- $CH_3-MgBr \xrightarrow[H_3O^{\oplus}]{CO_2} CH_3COOH$

Acetic acid

6. FROM ACYL HALIDES :-

$$R-\overset{O}{\overset{\|}{C}}-Cl + H-OH \longrightarrow R-\overset{O}{\overset{\|}{C}}-OH + HCl$$

Eg:- $CH_3-\overset{O}{\overset{\|}{C}}-Cl + H-OH \longrightarrow CH_3-\overset{O}{\overset{\|}{C}}-OH + HCl$

Ethanoyl chloride — Ethanoic acid

7. FROM ANHYDRIDES :-



$$R-\overset{O}{\overset{\|}{C}}-O-\overset{O}{\overset{\|}{C}}-R' + H_2O \longrightarrow RCOOH + R'COOH$$

Eg:- $C_6H_5-\overset{O}{\overset{\|}{C}}-O-\overset{O}{\overset{\|}{C}}-CH_3 + H_2O \longrightarrow C_6H_5COOH + CH_3COOH$

Benzoic ethanoic anhydride — Benzoic acid — Ethanoic acid

8. FROM ESTERS :-

$$R-\overset{O}{\overset{\|}{C}}-O-R' \underset{}{\overset{H_3O^{\oplus}}{\rightleftharpoons}} R-COOH + R'OH$$

Eg:-  $\overset{H_3O^{\oplus}}{\rightleftharpoons}$  $+ C_2H_5OH$

Ethyl benzoate — Benzoic acid

PHYSICAL PROPERTIES :-

BOILING POINT :-

Carbonylic acid have higher boiling point due to more extensive association of carbonylic acid molecules through intermolecular hydrogen bonding.

* The hydrogen bonds are not broken completely even in the vapour phase. Most carboxylic acid exist as dimer in vapour phase.

SOLUBILITY:-

* The solubility decreases with increase in number of carbon atoms.

* Higher carboxylic acids are practically insoluble in water due to the increased hydrophobic interaction of hydrocarbon part.

CHEMICAL REACTIONS:-

① REACTIONS INVOLVING CLEAVAGE OF O-H BOND:-

$$R-\overset{\overset{O}{\|}}{C}-O-H \longrightarrow RCOO^- + H^+$$

REACTION WITH Na:-

$$RCOOH + Na \longrightarrow RCOONa + \frac{1}{2} H_2O$$

REACTION WITH H_2O:-

$$RCOOH + H_2O \longrightarrow RCOO^- + H^+$$

ACIDITY:-

$$RCOOH + H_2O \longrightarrow RCOO^- + H^+$$

$$K_a = \frac{[RCOO^-][H^+]}{[RCOOH]}$$

$K_a \propto [H^+] \propto$ Acidity

$\Rightarrow pK_a \propto \frac{1}{K_a} \propto \frac{1}{\text{Acidity}}$

∵ Compounds which have more acidity value, their pK_a is less.


$$EDG-CH_2-\overset{\overset{O}{\|}}{C}-OH$$

$$EWG-CH_2-\overset{\overset{O}{\|}}{C}-OH$$


By putting EWG, OH bond polarity ↑ses. Hence, H^+ removal become easier. So EWG increases acidity.

② REACTIONS INVOLVING CLEAVAGE OF C-OH BOND:-

(i) FORMATION OF ANHYDRIDE :-



(ii) ESTERIFICATION :-



Ethyl ethanoate


(3) REACTIONS WITH PCl_5, PCl_3 and $SOCl_2$:-

$$RCOOH + PCl_5 \longrightarrow RCOCl + POCl_3 + HCl$$


$$3RCOOH + PCl_3 \longrightarrow 3RCOCl + H_3PO_3$$

$$RCOOH + SOCl_2 \longrightarrow RCOCl + SO_2 + HCl$$

(4) REACTION WITH AMMONIA :-



Eg :-



Ammonium pthalate


Phthalamide

$-NH_3$ Strong heating

Phthalimide

(3) REACTIONS INVOLVING -COOH GROUP :-

(i) REDUCTION :-




Eg :-

② DECARBOXYLATION :-

In this method, one C is decreased from its parent carbon.

$$R-\overset{O}{\overset{\|}{C}}-OH \xrightarrow[\Delta]{NaOH\ \&\ CaO} R-H + Na_2CO_3$$

Eg:- 

④ SUBSTITUTIONS REACTIONS IN THE HYDROCARBON PART :-


① HALOGENATION :-

HVZ REACTION :-
(Hell-Volhard-Zelinsky reaction)

$$R-CH_2-COOH \xrightarrow[(ii)\ H_2O]{(i)\ X_2/Red\ P} R-\underset{X}{CH}-COOH$$


Eg:- $CH_3-CH_2-COOH \xrightarrow[(ii)\ H_2O]{(i)\ Cl_2} CH_3-\underset{Cl}{CH}-COOH$

② ELECTROPHILIC SUBSTITUTION :-



COOH is an EWG. Hence it deactivates the benzene ring. Hence, next electrophile will attack at meta position.

(i) NITRATION :-



m-nitrobenzoic acid

(ii) BROMINATION :-

